


CELLULAR ENERGY COENZYME

NAD+

500 MG / 3 ML VIAL

1.67 mg PER UNIT



» KEY BENEFITS

- » Boosts cellular energy and ATP production
- » Activates sirtuins linked to longevity
- » Supports cognitive performance and focus
- » Enhances DNA repair and genomic stability
- » Improves mitochondrial function
- » Reduces fatigue and metabolic decline

CELLULAR ENERGY COENZYME

» NAD+

500 mg / 3 mL Vial

Reference scale showing the milligram dose delivered per unit on a U-100 insulin syringe (100 units = 1 mL). Pre-mixed in bacteriostatic water — no reconstitution required. Draw directly from the vial with a sterile insulin needle.

PER UNIT	TOTAL CONTENT	VOLUME	CONCENTRATION
1.67 mg	500 mg	3 mL	166.7 mg/mL

» UNIT ↔ MILLIGRAM REFERENCE

1 unit = 1.67 mg · U-100 insulin syringe (100 units = 1 mL)

UNITS	DOSE	UNITS	DOSE
1 unit	1.67 mg	12 units	20.00 mg
2 units	3.33 mg	15 units	25.00 mg
3 units	5.00 mg	20 units	33.33 mg
4 units	6.67 mg	25 units	41.67 mg
5 units	8.33 mg	30 units	50.00 mg
6 units	10.00 mg	40 units	66.67 mg
8 units	13.33 mg	50 units	83.33 mg
10 units	16.67 mg	100 units	166.67 mg

» HOW TO DOSE

- 01** Allow the vial to reach room temperature. Inspect the solution for clarity.
- 02** Wipe the rubber stopper with a sterile alcohol swab.
- 03** Using a sterile U-100 insulin syringe, draw the desired number of units directly from the vial.
- 04** Tap out air bubbles. Confirm the dose against the reference table above.
- 05** Inject subcutaneously. Rotate sites. Discard the syringe safely in a sharps container.